

Self-assess: Methods and Theory of Modern Optimisation

Andrew C. Eberhard Scott B. Lindstrom

January 10, 2022

Abstract

As necessary background for the course, you will need to have taken undergraduate courses in real analysis, basic vector calculus and linear algebra. It is also desirable to have some familiarity with a computer programming language like Matlab, Julia, Python, etc. The following questions will help you assess your readiness for the course.

1 Quiz

Vectors and matrices

Consider the matrices

$$A := \begin{vmatrix} 1 & 2 & 3 \\ 4 & 5 & 6 \\ 7 & 8 & 9 \end{vmatrix} \quad \text{and} \quad B := \begin{vmatrix} 1 & 2 \\ 3 & 4 \\ 5 & 6 \end{vmatrix}.$$

1. Show that A is invertible, or explain why it is not invertible.
2. Show that B is invertible, or explain why it is not invertible.
3. Calculate $A + B$ or explain why it does not exist.
4. Calculate AB or explain why AB does not exist.
5. Calculate BA or explain why BA does not exist.
6. Calculate $B^T A$ or explain why $B^T A$ does not exist.

Real analysis

7. Let $(X, \|\cdot\|)$ be a normed space. Suppose that the sequence x_n satisfies $\lim_{n \rightarrow \infty} x_n = x$ and also that the sequence y_n satisfies $\lim_{n \rightarrow \infty} \|y_n - x_n\| \rightarrow 0$. Show that $\lim_{n \rightarrow \infty} y_n = x$.

Vector calculus

8. Compute the gradient $\nabla f(x, y)$ where $f(x, y) = x^2 y + y e^{xy}$.

2 Solutions

1. A is invertible, because its column vectors are linearly independent. To verify this, suppose that $x_1v_1 + x_2v_2 + x_3v_3 = 0$ where v_1, v_2, v_3 are the column vectors of A from left to right and $x_1, x_2, x_3 \in \mathbb{R}$. This is just $Ax = 0$, and so we obtain the system

$$\left| \begin{array}{ccc|c} 1 & 2 & 3 & 0 \\ 4 & 5 & 6 & 0 \\ 7 & 8 & 10 & 0 \end{array} \right|$$

We can put this in reduced row echelon form as follows:

$$\left| \begin{array}{ccc|c} 1 & 2 & 3 & 0 \\ 0 & -3 & -6 & 0 \\ 0 & 0 & 1 & 0 \end{array} \right|,$$

which shows that $x = 0$.

2. B cannot be invertible, because it is not a square matrix.
3. $A + B$ does not exist because A and B do not have the same dimensions.
4. We calculate AB by matrix multiplication:

$$AB = \left| \begin{array}{cc} 1 \cdot 1 + 2 \cdot 3 + 3 \cdot 5 & 1 \cdot 2 + 2 \cdot 4 + 3 \cdot 6 \\ 4 \cdot 1 + 5 \cdot 3 + 6 \cdot 5 & 4 \cdot 2 + 5 \cdot 4 + 6 \cdot 6 \\ 7 \cdot 1 + 8 \cdot 3 + 10 \cdot 5 & 7 \cdot 2 + 8 \cdot 4 + 10 \cdot 6 \end{array} \right|.$$

5. BA does not exist because the number of rows of B does not match the number of columns in A .
6. $B^T A$ is computed as follows:

$$B^T A = \left| \begin{array}{ccc} 1 \cdot 1 + 3 \cdot 2 + 5 \cdot 3 & 1 \cdot 4 + 3 \cdot 5 + 5 \cdot 6 & 1 \cdot 7 + 3 \cdot 8 + 5 \cdot 10 \\ 2 \cdot 1 + 4 \cdot 2 + 6 \cdot 3 & 2 \cdot 4 + 4 \cdot 5 + 6 \cdot 6 & 2 \cdot 7 + 4 \cdot 8 + 6 \cdot 10 \end{array} \right|.$$

7. By the triangle inequality,

$$\|y_n - x\| \leq \|y_n - x_n\| + \|x_n - x\|.$$

Taking the limit as $n \rightarrow \infty$, we have that the right hand side approaches zero, which means that y_n converges to x .

8. The gradient consists of the partial derivatives in the variables x and y as follows:

$$\nabla f(x, y) = (2xy + y^2 e^{xy}, x^2 + yx e^{xy} + e^{xy})$$